

Pionic atoms probing πNN resonances

M. I. Krivoruchenko^{*,†)}, B. V. Martemyanov^{*,†)}, Amand Faessler^{†)} and C. Fuchs^{†)}

^{*}*Institute for Theoretical and Experimental Physics, B. Cheremushkinskaya 25
117259 Moscow, Russia*

^{†)}*Institut für Theoretische Physik, Universität Tübingen, Auf der Morgenstelle 14
D-72076 Tübingen, Germany*

The pion optical potential generated by the hypothetical πNN -coupled NN -decoupled dibaryon resonance $d'(2065)$ is calculated to the lowest order in nuclear matter density. The contribution to the pion optical potential is found to be within the empirical errors, so the $d'(2065)$ existence currently does not contradict to the observed properties of the π^- -nucleus bound states. Future progress in the pionic X-ray spectroscopy can reveal contributions of πNN resonances to energy levels and widths of the pionic atoms.

PACS numbers: 13.75-n, 13.75.Gx, 25.80.Hp, 14.20.Pt

The possibility for existence of dibaryon resonances is discussed over two decades [1, 2, 3, 4, 5]. The most popular candidates for exceptionally narrow dibaryon resonances are the H dihyperon with a mass below the $\Lambda\Lambda$ threshold, as predicted by Jaffe [1] within the framework of the MIT bag model, and the nonstrange πNN -coupled NN -decoupled d' dibaryon with quantum numbers $I = 0$ and $J^P = 0^-$ and a mass close to the πNN threshold [2, 3]. The resonance-like shape of the DCE cross section [6] was interpreted by Martemyanov and Schepkin [7] as a manifestation of the d' dibaryon with a mass of 2065 MeV. Searches of the H dihyperon did not give conclusive results [5, 8, 9, 10, 11]. The upper limits for the d' production cross sections are settled experimentally [12, 13], while the most specific features of the DCE reaction have been described using conventional mechanisms [14, 15]. Astrophysical constraints to masses and coupling constants of the dibaryons are derived from existence of massive neutron stars [16, 17, 18, 19].

The laboratory experiments with pionic atoms [20, 21, 22, 23, 24] provide one more opportunity to test existence of the d' dibaryon: The d' is coupled to the πNN channel, so as illustrated by Fig. 1 it contributes to the pion optical potential to the second order in the nuclear matter density and, as a result, manifests itself in shifts and broadening of the atomic spectral lines.

The pion-nucleus interaction effects on the pion bound states and present experimental status of the pionic X-ray spectroscopy are reviewed recently by Kienle and Yamazaki [25].

The d' contribution to the π^- optical potential at the nuclear density ρ_0 has the form

$$2\mu\delta V_{opt}(\mathbf{k}) = \frac{2g^2}{M\eta_F} \int \frac{d\mathbf{p}_1}{(2\pi)^3} \frac{d\mathbf{p}_2}{(2\pi)^3} \frac{1}{|D_J|^2} \frac{\theta(p_F - |\mathbf{p}_1|)\theta(p_F - |\mathbf{p}_2|)}{\mathbf{p}_1^2/(2m) + \mathbf{p}_2^2/(2m) + \mathbf{k}^2/(2\mu) - (\mathbf{p}_1 + \mathbf{p}_2 + \mathbf{k})^2/(2M) - \Delta M + i\Gamma/2} \quad (1)$$

where $\mathbf{p}_1$ and $\mathbf{p}_2$ are momenta of two protons, $\mathbf{k}$ is the pion momentum, m is the nucleon mass, μ is the pion mass, p_F is the Fermi momentum, $\Delta M = M - 2m - \mu$, $M = 2065$ MeV and Γ are the d' mass and width, $g = 9.2 \text{ GeV}^{-1}$ is the $\pi NN d'$ coupling constant determined from the width $\Gamma = 0.5 \text{ MeV}$ of d' decay in the vacuum [6, 26]

$$\Gamma = \frac{3g^2}{64\pi^2} \sqrt{\mu M} \Delta M^2 \quad (2)$$




FIG. 1: Contribution to the pion optical potential of the d' dibaryon (solid double line). The single solid lines show the proton holes (p_h) in the Fermi sphere, the dashed lines show the pion propagating in the nuclear matter.

TABLE I: Contribution of the d' dibaryon to the parameters B_0 and C_0 of the pion optical potential in the symmetric nuclear matter at the saturation density $\rho_0 = 0.17 \text{ fm}^{-3}$ and an effective density $\rho_e \sim 0.6\rho_0$ where the pion-nucleus interaction has its maximum, as compared to the empirical values extracted from the pionic X-ray spectroscopy using the standard (S) model, Weise-Friedman (WF) model [35, 36], and Kienle-Yamazaki (KYa) model [25]. The empirical data are from [37] and [25], respectively. The values in parenthesis refer to the case of in-medium width of d' .

Model	$\mu^4 \text{Re}B_0$	$\mu^4 \text{Im}B_0$	$\mu^6 \text{Re}C_0$	$\mu^6 \text{Im}C_0$
S	-0.15 ± 0.04	0.054 ± 0.002	-0.28 ± 0.01	0.062 ± 0.003
WF	-0.07 ± 0.04	0.053 ± 0.002	-0.28 ± 0.01	0.067 ± 0.003
KYa	-0.02 ± 0.02	0.047 ± 0.002		
$d'[\rho_0]$	0.004	0.002	0.004	0.010
$d'[\rho_e]$	0.005	0.001	0.013	0.008

with enhancement factor η_F included to the coupling constant g^2 . The Jost function D_J in the effective-range approximation is given by [27]

$$D_J = \frac{|\mathbf{p}_1 - \mathbf{p}_2|/2 + i\gamma}{|\mathbf{p}_1 - \mathbf{p}_2|/2 + i\alpha} \quad (3)$$

where $\alpha \approx 2/r_e$ and $\gamma = 2/a/\alpha/r_e$, with $a = 7.82 \text{ fm}$ being the effective proton-proton scattering length including the Coulomb interaction [29], and $r_e = 2.67 \text{ fm}$ the effective radius. The enhancement factor $\eta_F \approx 1 + 32/(r_e^2 M \Delta M) \approx 3$ according to [28]. It is determined as an average value of the $1/|D_J|^2$ over the three-body πNN phase space.

The d' in-medium width is estimated to be $\Gamma^* = 10 \text{ MeV}$ [6]. It appears due to the reactions $d'N \rightarrow NNN$. It is of the first order in the density, so formally we go beyond the lowest second-order calculation. The effect, however, is numerically large. The d' width is increased as compared to the vacuum value by a factor of 20. The vacuum decay channel $d' \rightarrow \pi NN$ is blocked in our case, since the pion is bound. The collision broadening of the d' is the only effect contributing to the imaginary parts of the d' optical potential.

The second-order pion optical potential is related to the non-resonant absorption of pions on nucleon pairs [30]. It is parameterized in the form

$$2\mu\delta V_{opt}(\mathbf{k}) = \delta q + \delta p \mathbf{k}^2 \quad (4)$$

where

$$\delta q = -16\pi(1 + \frac{\mu}{2m})B_0\rho_p\rho_n, \quad (5)$$

$$\delta p = -16\pi(1 + \frac{\mu}{2m})^{-1}C_0\rho_p\rho_n \quad (6)$$

are the s - and p -wave parts of the pion optical potential, ρ_p and ρ_n are the proton and neutron densities. Eq.(1) determines the d' contribution to the parameters B_0 and C_0 .

The results are summarized and compared to the experiment in Table 1. The empirical data are taken from [37] for two realistic models with a vanishing nucleon-nucleon correlations parameter. The results of fit of the s -wave part of the π^- - nucleus potential [25] are also given. The real s -wave part should be of the same magnitude as the imaginary part [31, 32, 33, 34]. The result of the standard model $|\text{Re}C_0| \sim 3|\text{Im}C_0|$ is not fully satisfactory. The phenomenological pion-nucleon isovector scattering length in the standard model also appears to be overestimated as compared to the free value [25, 37]. A specific version of the relativistic impulse approximation [36, 38, 39, 40, 41, 42] and the recent WF model [35, 36] reconcile those problems, at least partially.

Due to a strong repulsive π^- -nucleus interaction and an attractive Coulomb interaction, the pions are bound at the surface of nuclei at an effective density $\rho_e \sim 0.6\rho_0$ [25]. The nontrivial momentum dependence of the Jost function and the Breit-Wigner amplitude results in a density dependence of the parameters B_0 and C_0 . Table 1 shows, respectively, two sets of the d' parameters for ρ_0 and ρ_e . The in-medium d' width is scaled according to the reduced density.

The $d'(2065)$ corrections to the B_0 and C_0 at $\rho_e \sim 0.6\rho_0$ where the pion-nucleus interaction has the strongest effect on the pionic atoms are currently within the empirical errors. If πNN -coupled resonances exist, the progress in pionic X-ray spectroscopy can reveal their contributions to energy levels and widths of the π^- -nucleus bound states.

The authors are grateful to M. G. Schepkin for useful remarks. M.I.K. and B.V.M. wish to acknowledge kind

hospitality at the University of Tuebingen. This work is supported by DFG grant No. 436 RUS 113/721/0-1 and RFBR grant No. 03-02-04004.

-
- [1] R. L. Jaffe, Phys. Rev. Lett. **38**, 195 (1977); Erratum: Ibid **38**, 617 (1977).
 - [2] P. J. Mulders, A. T. Aerts and J. J. de Swart, Phys. Rev. **D21**, 2653 (1980).
 - [3] L. A. Kondratyuk, B. V. Martemyanov and M. G. Schepkin, Sov. J. Nucl. Phys. **45**, 776 (1987).
 - [4] L.V. Fil'kov, V.L. Kashevarov, E.S. Konobeevski, M.V. Mordovskoy, S.I. Potashev, V.M. Skorkin, Phys. Rev. **C61**, 044004 (2000).
 - [5] T. Sakai, K. Shimizu, K. Yazaki, Prog. Theor. Phys. Suppl. **137**, 121 (2000).
 - [6] R. Bilger, H. A. Clement and M. G. Schepkin, Phys. Rev. Lett. **71**, 42 (1993).
 - [7] B. V. Martemyanov and M. G. Schepkin, JETP Lett. **53**, 139 (1991).
 - [8] S. Aoki et al., Phys. Rev. Lett. **65**, 1729 (1990).
 - [9] V. V. Barmin et al., Phys. Lett. **B370**, 233 (1996).
 - [10] J. Belz et al., Phys. Rev. Lett. **76**, 3277 (1996); Phys. Rev. **C56**, 1164 (1997).
 - [11] R. E. Chrien, Nucl. Phys. **A629**, 388c (1998).
 - [12] U. Siodlaczek et al., Eur. Phys. J. **A9**, 309 (2000).
 - [13] E. Doroshkevich et al., Eur. Phys. J. **A18**, 171 (2003).
 - [14] M. Nuseirat, M.A.K. Lodhi, M.O. El-Ghossain, W.R. Gibbs, W.B. Kaufmann, Phys. Rev. **C58**, 2292 (1998).
 - [15] Yu-X. Liu, Amand Faessler, J. Schwieger and A. Bobyk, J. Phys. **G24**, 1135 (1998).
 - [16] M. I. Krivoruchenko, JETP Lett. **46**, 3 (1987).
 - [17] A. Faessler, A. J. Buchmann and M. I. Krivoruchenko, Phys. Rev. **C56**, 1576 (1997).
 - [18] A. Faessler, A. J. Buchmann, M. I. Krivoruchenko and B. V. Martemyanov, J. Phys. **G24**, 791 (1998).
 - [19] N. K. Glendenning and J. Schaffner-Bielich, Phys. Rev. **C58**, 1298 (1998).
 - [20] J. Koni et al., Nucl. Phys. **A326**, 401 (1979).
 - [21] A. Olin et al., Nucl. Phys. **A439**, 589 (1985).
 - [22] C. T. A. M. de Laat et al., Phys. Lett. **B162**, 81 (1985).
 - [23] T. Yamazaki et al., Phys. Lett. **B418**, 246 (1998).
 - [24] H. Geissel et al., Phys. Rev. Lett. **88**, 122301 (2002).
 - [25] P. Kienle and T. Yamazaki, Prog. Part. Nucl. Phys. **52**, 85 (2004).
 - [26] M. Shchepkin, O. Zaboronsky, H. Clement, Z. Phys. **A345** 407 (1993).
 - [27] M. L. Goldberger and K. M. Watson, *Collision Theory*, John Wiley and Sons, NY (1965).
 - [28] S. M. Kiselev, M. I. Krivoruchenko, B. V. Martemyanov, Amand Faessler and C. Fuchs, Nucl. Phys. **A650**, 78 (1999).
 - [29] G. E. Brown and A. D. Jackson, *The Nucleon-Nucleon Interaction*, North-Holland Publishing Company, Amsterdam, 1976.
 - [30] M. Ericson and T. E. O. Ericson, Ann. Phys. (N.Y.) **36**, 323 (1966).
 - [31] C. Garcia-Recio, E. Oset and L. L. Salcedo, Phys. Rev. **C37**, 194 (1988).
 - [32] E. Oset, C. Garcia-Reico and J. Nieves, Nucl. Phys. **A584**, 653 (1995).
 - [33] C. M. Ko and D. O. Riska, Nucl. Phys. **A312**, 217 (1978).
 - [34] L. L. Salcedo, K. Holinde, E. Oset and C. Schutz, Phys. Lett. **B353**, 1 (1995).
 - [35] W. Weise, Nucl. Phys. **A690**, 98 (2001).
 - [36] E. Friedman, Phys. Lett. **B524**, 87 (2002).
 - [37] E. Friedman, Nucl. Phys. **A710**, 117 (2002).
 - [38] B. L. Birbrair, V. N. Fomenko, A. B. Gridnev and Yu. A. Kalashnikov, J. Phys. **G9**, 1473 (1983); **G11**, 471 (1985).
 - [39] P. F. A. Goudsmit, H. J. Leisi and E. Matsinos, Phys. Lett. **B271**, 290 (1991).
 - [40] B. L. Birbrair and A. B. Gridnev, Nucl. Phys. **A528**, 647 (1991).
 - [41] A. Gal, B. K. Jennings and E. Friedman, Phys. Lett. **B281**, 11 (1992).
 - [42] B. L. Birbrair, A. B. Gridnev, L. P. Lapina, A. A. Petrunin and A. I. Smirnov, Nucl. Phys. **A547**, 645 (1992).